



international linear collider

Рождение пар скалярных топ-кварков на будущем линейном ускорителе ИЛС и оценка основного фонового процесса

A.Bartl (**Univ. of Vienna**)


W.Majerotto **HEPHY (Vienna)**

K.Moenig **DESY (Zeuthen)**

A.N.Skachkova, N.B.Skachkov

JINR (Dubna)

Experimental restrictions on the STOP mass



“ Search for the pair production of scalar top quarks in the acoplanar charm jet final state in $p\bar{p}$ collisions at $\sqrt{S} = 1.96$ TeV”

D0 Note 5134-CONF

7 June 2006


Fermilab-Pub-06/396-E

hep-ex/0611003

Phys. Lett. B645 (2007)

119-127

Simulation is done by use of PYTHIA 6.4 + CIRCE 1




$e^+ e^-$ CM energy = 500 GeV

STOP pair production
cross section

$$\sigma = 2.37 \text{ fb}$$

TOP pair production
cross section

$$\sigma = 35.9 \text{ fb}$$



In order to simulate the STOP pair production, we assumed the following scenario for the MSSM model parameters:

- $M_{\tilde{Q}} = M_{\tilde{t}_L} = 270 \text{ GeV}$ (left squark mass)
- $M_{\tilde{U}} = M_{\tilde{t}_R} = 270 \text{ GeV}$ (right squark mass)
- $A_t = -500 \text{ GeV}$ (top and bottom trilinear coupling)
- $\mu = -370 \text{ GeV}$
- $\tan\beta = 5$
- $M_1 = 80 \text{ GeV}$
- $M_2 = 160 \text{ GeV}$

Corresponds to

$$M_{\text{stop}} = 167.9 \text{ GeV,}$$

$$M_{\chi_1^0} = 80.9 \text{ GeV}$$

$$M_{\chi_{1\pm}} = 159.2 \text{ GeV}$$

Our aim is:


- *To find out physical variables (Energy, PT, angle and invariant mass distributions) most suitable for signal (stop) / background (top) separation*
- *To estimate the corresponding values of cuts on these variables*

Cross section dependence on E_{beam}


(without any cuts)

$2E_{\text{beam}}[\text{GeV}]$	$\sigma_{\text{stop}} [\text{fb}]$	N_{stop}	$\sigma_{\text{top}} [\text{fb}]$	N_{top}
350	0.23	233	13.76	13750
400	1.34	1347	38.79	38740
<u>500</u>	<u>2.37</u>	<u>2378</u>	<u>35.94</u>	<u>35950</u>
800	1.89	1809	17.36	17359
1000	1.42	1265	11.66	11656

e^+ , e^- beam energy spectrum from CIRCE 1



Electron e^- (positron e^+) beam energy with account of beamstrahlung



Correlation between e^+ and e^- beam spectra





$$Y_i = E^i / E_{\text{beam}}^i \quad (i = e^+, e^-)$$

The peak luminosity is supposed to be $2 \cdot 10^{34} \text{cm}^{-2}\text{s}^{-1}$.

The total luminosity required is 1000fb^{-1} during the first phase of operation at

$$2E_{\text{beam}}^e = \sqrt{S_{ee}} = 500 \text{ GeV}.$$

Main Scalar top quark distributions





Invariant mass of 4 jets and M_{missing} variable



Good for Signal / Background separation cut

$$M_{\text{inv}}(4 \text{ jets}) < 160 \text{ GeV} \quad \text{and} \quad M_{\text{missing}} > 250 \text{ GeV}$$

STOP



TOP



$M_{\text{inv}}(4 \text{ jets})$

M_{missing}

Used cuts for S/B separation

- 1.) The events with clear recognized 2 B-jets (according to PYTHIA)
(B-jet is determined as a jet that includes b-meson)

Stop cut efficiency = 0.84

Top cut efficiency = 0.94

But, in the experiment only 50% efficiency of the B-jets and B_{bar} -jets separation and the 80% of the corresponding purity is expected

- 2.) Invariant mass of 4 jets (b_{jet} , $b_{\text{bar}_{\text{jet}}}$, 2jets_W) M_{inv} (All jets) < 160 GeV
together with the cut above

Stop cut efficiency = 0.78

Top cut efficiency = 0.001

- 3.) Invariant Missing mass $M_{\text{miss}} > 250$ GeV
together with the cuts above

Stop cut efficiency = 0.94

Top cut efficiency = $6 \cdot 10^{-6}$

Achieved S/B ratio = 143





*The rest is only **13 background events per year**, while for the Signal events – **1086/year** (for the integrated Luminosity $L=1000 \text{ fb}^{-1}/\text{year}$)*

Cross section dependence on E_{beam}

(with the cuts above)

$2E_{\text{beam}}[\text{GeV}]$	$\sigma_{\text{stop}} [\text{fb}]$	N_{stop}	$\sigma_{\text{top}} [\text{fb}]$	N_{top}
350	0.0089	8	0	0
400	0.52	521	$2.32 * 10^{-4}$	0.2
<u>500</u>	<u>1.80</u>	<u>1806</u>	<u>$2.26 * 10^{-2}$</u>	<u>12.6</u>
800	0.99	995	$1.08 * 10^{-2}$	10
1000	0.41	410	$6.26 * 10^{-3}$	6

μ distributions in the signal events




Signal μ 's

Fake μ 's


Invariant mass of b_{jet} & $2jets_w$

Good for Signal / Background separation cut $M_{inv}(b_{jet}, 2jets_w) < 100$ GeV!

STOP



TOP





$Minv(b\text{-jet}, 2jets_w)$

The most important variable - invariant mass of b_{jet} & $2jets_W$

In the case of **TOP pair** production
it gives

The reconstruction of M_{Top} (175 GeV):

$$M_{inv}(B_{jet} \& 2jets_W) = M_{Top}$$



Stop invariant mass



The reconstruction of the STOP
invariant mass M_{STOP} (167.9 GeV):

$$M_{inv}(STOP) =$$

$$M_{\chi_1^0} + M_{inv}(b_{jet}, 2jets_W) =$$

$$= M_{\chi_1^0} + \sqrt{(P_{b_{jet}} + P_{jet1_W} + P_{jet2_W})}$$

But χ_1^0 - is not detectable particle,
its mass should be known from the
other experiments



Invariant mass of b_{jet} & $2jets_W$ gives

For the case of



STOP pair production

$$M_{inv}(B_{jet}, 2jets_W) = M_{inv}(STOP) - M_{\chi_1^0}$$

The peak of $M_{inv}(b_{jet}, 2jets_W) \approx 87 \text{ GeV}$


$$M_{\chi_1^0} \approx 80.9 \text{ GeV}$$

$$\begin{aligned} M_{stop} &= M_{\chi_1^0} + M_{inv}(b_{jet}, 2jets_W) = \\ &= 167.9 \text{ GeV} \end{aligned}$$



The test of the other Scalar top mass

- $M_{stop} = 200.1 \text{ GeV}$
- $M_{\chi_1^0} = 80.9 \text{ GeV}$
- $M_{\chi_1^+} = 159.6 \text{ GeV}$



Right edge of the peak of

$$M_{inv}(b_{jet}, 2jet_W) \approx 120 \text{ GeV}$$

$$M_{\chi_1^0} \approx 80 \text{ GeV}$$

$$M_{stop} = M_{\chi_1^0} + M_{inv}(b_{jet}, 2jet_W) = 200 \text{ GeV}$$

509 events
S/B = 40

Conclusion

1. *The MC (PYTHIA 6.4 + CIRCE 1) study of stop pair production in e^+e^- collisions was done at $\sqrt{S}_{ee} = 350, 400, 500, 800, 1000$ GeV.*
2. *The detailed analysis done at $\sqrt{S}_{ee} = 500$ GeV has shown that proposed 3 cuts allow to reach $S/B = 143$.*
3. *A possibility of a good reconstruction of the M_{STOP} from the peak position of M_{inv} (3 jets, i.e. $b_{jet} + 2 jets_W$) distribution is demonstrated.*

So, finally, the channel

$$STOP STOP \rightarrow b \chi_1^+ b \chi_1^- \rightarrow b \bar{b} q \bar{q}' \mu^+ \nu_\mu \chi_1^0 \chi_1^0$$

is VERY promising for STOP quark search!



The subsequent decay channels have been considered:

➤ $STOP\ STOP \rightarrow b\ \chi_1^+\ b_{bar}\ \chi_1^- \rightarrow b\ b_{bar}\ q_i\ q_j\ bar\ \mu^-\ \nu_\mu\ \chi_1^0\ \chi_1^0$

➤ $t\ t \rightarrow b\ W^+\ b_{bar}\ W^- \rightarrow b\ b_{bar}\ q_i\ q_j\ bar\ \mu^-\ \nu_\mu$

The only difference of STOP / TOP production is the presence of the two non-detectable neutralinos in the case of stop pair production.

The quarks hadronize into jets. Jets are determined by use of PYCLUS jetfinder based on “Durham” cluster distance measure algorithm.

Both the signal and background events have the same experimental signature (b & b_{bar} - jets, 2 jets from $W \rightarrow q_i\ q_j$ decay and μ^-).

Publications & previous presentations

- *“Pair production of scalar top quarks in e^+e^- collisions at ILC.”*

Authors: A.Bartl, W.Majerotto, K.Möniq, A.N.Skachkova, N.B.Skachkov

arXiv: 0804.2125,

ILC-NOTE-2008-042

- *“Pair production of scalar top quarks in e^+e^- collisions at ILC.”*

Talk at the “QUARKS-2008” (<http://quarks.inr.ac.ru/>)

15th International Seminar on High Energy Physics

Sergiev Posad, Russia, 23-29 May, 2008.

To be published in the proceedings

Publications & previous presentations

- *“Stop pair production in polarized photon-photon collisions”.*

A.Bartl, W.Majerotto, K.Moenig, A.Skachkova, N.Skachkov.

Talk at the “International Conference on Linear Colliders”, LCWS04, 19-23 April 2004, Paris, France
Proceedings of the “International Conference on Linear Colliders”; LCWS 04, volume II, p.919-922

- *“Stop pair production at ILC”* (<http://www.linearcollider.org/cms/?pid=1000364>)

Talk at the International Linear Collider Workshop (14-17 November 2005, Vienna, Austria)

- *“Stop pair production in photon-photon collisions at ILC”*

Десятая научной конференции молодых ученых и специалистов ОИЯИ. Дубна, 6-10 февраля 2006 г.
Труды десятой научной конференции молодых ученых и специалистов ОИЯИ. Дубна, 6-10 февраля 2006 г. стр. 135-138, ISBN 5-9751-0024-0



- *“Pair production of scalar top quarks in polarized photon-photon collisions at ILC.”*

Authors: A.Bartl, W.Majerotto, K.Mönig, A.N.Skachkova, N.B.Skachkov



arXiv: **0804.1700**, **ILC-NOTE-2007-036**

B-quarks distributions

STOP





TOP





B-jets distributions

STOP





TOP





W-quarks distributions

STOP





TOP





Distributions of jets from W decay

STOP





TOP





Missing energy (ν_μ , $\sim\chi_1^0$, beam pipe) and detected energy distributions

Good for Signal / Background separation with cut $E_{cal_tot} < 180$ GeV

STOP



TOP



Missing energy



Detected energy

Total scalar Σ PT and Invariant mass of 4jets + μ



Good for Signal / Background separation with the cuts

$PT_{skalsum} < 150 \text{ GeV}$ and $M_{inv} (4 \text{ jets} + \mu) < 200 \text{ GeV}$!

STOP



TOP



Scalar Σ PT

$M_{inv} (4 \text{ jets} + \mu)$